

# Quasi-Two-Dimensional Modeling of the Self Gravitating Gas \*

A. Gromov<sup>†</sup>, V. Perepelovsky<sup>‡</sup>

## Abstract

The quasi-two-dimensional modeling of the small adiabatic perturbation on the background of the stationary configuration of the selfgravitating gas with the weak transverse nonhomogeneity approximation is presented. The space periodic character of the solution of this system is proofed.

## 1 The Physical Model

An effects arising during an expansion of the plane stationary adiabatic wave with a small amplitude on the background of the stationary configuration of the self gravitating gas have a volume nonlinear character. These effects arising in the process of competition of the two basic mechanisms: the self gravitating and thermal expansion of particles. Three physical parameters define the solution of this problem:  $\Phi_0$  - the characteristic gravitational potential,  $T_0$  - the characteristic temperature for the thermal expansion and  $\gamma$

---

\*Cosmoparticle physics. 1. Proceedings of 1 International conference on cosmoparticle physics "Cosmion-94", dedicated to 80 Anniversary of Ya.B.Zedovich and 5 Memorial of A.D.Sakharov. Eds M.Yu.Khlopov, M.E.Prokhorov, A.A.Starobinsky, J.Tran Thanh Van, Editions Frontieres, 1996. p.p.203 - 206.

<sup>†</sup>Computer Science Chair, Faculty of Technical Cybernetics, St. Petersburg State Technique University, Dom 29, Politechnitheskaja Ulitsa, 192251, St. Petersburg, RUSSIA, E-mail gromov@natus.stud.pu.ru

<sup>‡</sup>Department of Electron-Ion Processing of Solids, St.Petersburg Electrotechnical University, Prof. Popov str.,5 St.Petersburg, 197376, Russia, FAX +7-(812)-234-9983, E-mail borisk@sovam.com

- the index of adiabatic. The stationary configuration of the self gravitating gas has cylindrical symmetry [1]. We use the term "radius" to fix the boundary between the areas of the gas and vacuum. The wave under consideration perturbed to a small extent the initial distribution of the speed, density and radius. Smallness of the perturbation is of principle character. The mathematical description of this model is represented by the system of equation of motion and continuity (three dimensional nonstationary partial differential equations), Poisson equation and algebraic equation of state:

$$\frac{\partial \vec{v}}{\partial t} + (\vec{v} \cdot \nabla) \vec{v} = -\frac{\nabla P}{\rho} - \nabla \Phi; \quad (1)$$

$$\frac{\partial \rho}{\partial t} + \nabla \cdot (\rho \vec{v}) = 0; \quad (2)$$

$$\nabla^2 \Phi = 4\pi G \rho; \quad (3)$$

$$P = A \rho^\gamma, \quad (4)$$

there  $P$  is pressure,  $\vec{v}$  - thermal speed,  $\Phi$  - gravitational potential,  $\rho$  - gas density,  $t$  - time,  $A = \text{const.}$  This system describes the ideal classic gas with self gravitation. In the system of coordinate related to the stationary wave the derivative of time disappear [1] from the system (4). An addition to (4) we'll use also the radius component of the equation of motion multiplyed by  $\rho(z, r)r$ :

$$\frac{1}{2} \rho r \frac{\partial v_r^2}{\partial r} + r \frac{\partial P}{\partial r} = -\rho r \frac{\partial \Phi}{\partial r}. \quad (5)$$

## 2 The Relations of the Quasi-Two-Dimensional Averaging Method

The QTDA method is applied to finite three dimensional space problem with separate direction. A rigorous and consistent quasi-two-dimensional approach must based on the cross-section averaging of three dimensional physical model equations. This procedure allows one to proceed from three-dimensional distributions of physical quantities (such as potential, mass density etc.) to their one-dimensional averaged distributions [2]. The following

relations of the QTDA method have been used:

$$\int_{S(z)} \nabla^2 \Phi(r, z) ds = \frac{d}{dz} \left\{ \left( \frac{d\Phi(z)}{dz} \right)^S S(z) \right\} + \oint_{L(z)} \frac{\vec{n} \cdot \nabla \Phi(r, z)}{\cos \vartheta} dl, \quad (6)$$

$$\int_{S(z)} \nabla \Phi(r, z) \cdot d\vec{s} = \frac{d}{dz} \left\{ \left( \Phi(z) \right)^S S(z) \right\} + \oint_{L(z)} \frac{\vec{n} \cdot \vec{e}_z \Phi(r, z)}{\cos \vartheta} dl, \quad (7)$$

$$\int_{S(z)} \nabla \cdot \vec{A}(r, z) ds = \frac{d}{dz} \left\{ \left( A_z(z) \right)^S S(z) \right\} + \oint_{L(z)} \frac{\vec{n} \cdot \vec{A}(r, z)}{\cos \vartheta} dl, \quad (8)$$

where  $\vec{n}$  - external (as referred to the section  $S(z)$ ) normal to the side surface;  $\vartheta$  is the angle between the axis  $r$  and the normal  $\vec{n}$  to the side surface of the cylinder;

$$\left( \frac{d\Phi(r, z)}{dz} \right)^S S(z) = \int_{S(z)} \nabla \Phi(r, z) \cdot d\vec{s}; \quad (9)$$

$$(\Phi(z))^S S(z) = \int_{S(z)} \Phi(r, z) ds, \quad (10)$$

where index  $^S$  marks the cross section averaging values. All characteristics of the three dimensional problem are described in the new equations by the relation establishment between new functions - the average quantity variable of physical value and the boundary meanings of the initial function. In this formulas the boundary meanings are represented by the contour integrals. It is seen that all non-one dimensional effects are located in the contour integrals.

### 3 The QTD Model of the Weak Transverse Nonhomogeneity Approximation

The QTDA equations are the result of application of the QTDA method represented by (3) - (10) to equations of continuity, Poisson, and (2) in the system of coordinate related to the wave. In this model we studying the effects related to the small perturbations of the physical values. According to this

the smallness of following parameters have been supposed:

$$\mu = -\frac{v_r^2(R) - (v_r^2)^S}{c_0^2}, \quad (11)$$

$$d = \frac{(\rho^\gamma)^S - (\rho^S)^\gamma}{\rho_0^\gamma}, \quad (12)$$

where  $c_0$  and  $\rho_0$  are the sound speed and density on the axis  $z$  in the nonperturbated part of the cylinder. The smallness of these parameters correspond to the weak transverse nonhomogeneity approximation. We study the equations of continuity, Poisson, state and (2) in the 0-approximation of small parameters (12). The structure of the QTDA equations depended from contour integrals, which are simplified in the 0-approximation because in this case: 1)  $\cos \theta = 1$ ; 2)  $\vec{n}$  is parallel to  $\nabla \Phi$ ;  $\vec{n} \cdot \vec{e}_z = 0$ . Therefore only the Poisson equation has a nontrivial summand after the averaging. Thus the QTDA equations under consideration become as follows:

$$\frac{1}{2}(v^2)^S = -\frac{A\gamma}{\gamma - 1}(\rho^S)^{\gamma-1} - \Phi^S + C, \quad (13)$$

where  $C = const$ ; equation of continuity:  $\rho^S v^S = const$ ; Poisson equation:

$$\frac{d^2 \Phi^S(z)}{dz^2} - \frac{2}{R^2} \Phi^S(z) = 4\pi G \rho^S(z). \quad (14)$$

It is not possible to obtain the second summand from one dimensional model. The averaging of the 0-approximation of the equation (2) represent the dependence of the radius of the cylinder from the coordinate  $z$ :

$$\frac{dR}{dt} = v_r(R) = v_z K(\sigma) \frac{d\sigma}{dz}, \quad (15)$$

where  $\sigma$  is the surface density,  $K(\sigma) = \frac{2-\gamma}{2(1-\gamma)} \sigma^{\frac{2-\gamma}{2(1-\gamma)}-1}$ . Let's estimate benefit of transverse motion energy in the full energy of the system. For this purpose we'll substitute  $\vec{v} = \vec{v}_r + \vec{v}_z$  in (13) and compare the energy of longitudinal and transverse motions we find out, that the latters are unessential far from the wave front  $\Delta z$ ,

$$\Delta z \gg \max \left| \frac{v_z(R) K(\sigma)}{\sqrt{F_i}} \Delta \sigma \right|, \quad (16)$$

where  $F_i$  - one of the set from  $\left((v_z^2)^S, \frac{2A\gamma}{\gamma-1}(\rho^S)^{\gamma-1}, 2|\Phi^S|, 2|C|\right)$ .  $\Delta\sigma$  - is a changing of surface density along the length  $\Delta z$ . It means that volume effects are more valuable than valuable surface ones. We'll use in following the averaging values without index  $S$ .

## 4 The QTD Description of the Self Gravitating Gas

For future investigation the dimensionless variables (the coordinate, density, pressure and potential) have been introduce:

$$\xi = \frac{r}{L_j}, \quad \delta = \frac{\rho}{\rho_0}, \quad p = \frac{P}{P_0}, \quad \varphi = \frac{\Phi}{\Phi_0}, \quad (17)$$

$$L_j^2 = \frac{\pi c_0^2}{G\rho_0}, \quad c_0^2 = \left(\frac{dp}{d\rho}\right)_0 = A\gamma\rho_0^{\gamma-1}, \quad (18)$$

where  $L_j$  - is the Djeence's length,  $c_0$  - sound speed. There are characteristic values marked with index "0" in this formulas. The dimensionless equations are:

$$\dot{\varphi} = u(1 - \gamma\delta^{\gamma+1}), \quad (19)$$

$$\dot{u} = (\varphi + \delta)(1 - \gamma\delta^{\gamma+1}), \quad (20)$$

$$\dot{\delta} = \beta u \delta^3, \quad (21)$$

where  $\dot{\phantom{x}} = \frac{d}{d\tau}$ ,  $d\xi = (1 - \gamma)\delta^{\gamma+1}d\tau$ . The meaning of density related to sound speed is  $\delta_s = \left(\frac{1}{\gamma}\right)^{\frac{1}{\gamma+1}}$ , so the (21) is not equal to infinity. The first integral of (21) is

$$\varphi = g(\delta) + C, \quad (22)$$

$$g(\delta) = -\frac{1}{\beta} \left( \frac{1}{2\delta^2} + \frac{\gamma}{\gamma+1} \delta^{\gamma-1} \right), \quad (23)$$

where  $C = \text{const}$ . From the system of equations

$$\frac{du}{d\tau} = (1 - \gamma\delta^{\gamma+1})(g(\delta) + \delta + C), \quad (24)$$

$$\frac{d\delta}{d\tau} = \beta u \delta^3, \quad (25)$$

the second integral of the system (25) has been received and has a form of "energy conservation law":  $E = \frac{u^2}{2} + U$ , where  $u$  plays a role of speed,  $E$  is the constant "energy", and the "potential"

$$U = -\frac{1}{2}[C + g(\delta)]^2 + \frac{1}{\beta}(\frac{1}{\delta} + \delta^\gamma). \quad (26)$$

The local minimum of the potential  $U$  correspond to finite solutions of the system (25). It is according to result have received in [1] for  $\gamma = 2$ .

## 5 Acknowledgements

I'm grateful to Prof. Arthur D. Chernin for encouragement and discussion. This paper was financially supported by "COSMION" Ltd., Moscow.

## References

- [1] V.A.Antonov, A.D.Chernin, Non-Linear Waves in a Gravitating Gas, Sov.Astronomy, **54** 2 p.315, 1977 (in Russian).
- [2] A.A.Barubin, V.V.Perepelovsky, Method of transverse averaging for quasi-two-dimensional model of thin film semiconductor structures, Electronic Engineering, Ser. 3, Microelectronics, 1991, No.3(142), pp.12-15 (in Russian).